



chromosome 9

Humans normally have 46 chromosomes in each cell, divided into 23 pairs. Two copies of chromosome 9, one copy inherited from each parent, form one of the pairs. Chromosome 9 is made up of about 141 million DNA building blocks (base pairs) and represents approximately 4.5 percent of the total DNA in cells.

Identifying genes on each chromosome is an active area of genetic research. Because researchers use different approaches to predict the number of genes on each chromosome, the estimated number of genes varies. Chromosome 9 likely contains 800 to 900 genes that provide instructions for making proteins. These proteins perform a variety of different roles in the body.

Health Conditions Related to Chromosomal Changes

The following chromosomal conditions are associated with changes in the structure or number of copies of chromosome 9.

9q22.3 microdeletion

9q22.3 microdeletion is a chromosomal change in which a small piece of the long (q) arm of chromosome 9 is deleted in each cell. Affected individuals are missing at least 352,000 base pairs, also written as 352 kilobases (kb), in the q22.3 region of chromosome 9. This 352-kb segment is known as the minimum critical region because it is the smallest deletion that has been found to cause the signs and symptoms related to 9q22.3 microdeletions. These signs and symptoms include delayed development, intellectual disability, certain physical abnormalities, and the characteristic features of a genetic condition called Gorlin syndrome (also known as nevoid basal cell carcinoma syndrome). 9q22.3 microdeletions can also be much larger; the largest reported deletion included 20.5 million base pairs (20.5 Mb).

People with a 9q22.3 microdeletion are missing two to more than 270 genes on chromosome 9. All known 9q22.3 microdeletions include the *PTCH1* gene. Researchers believe that many of the features associated with 9q22.3 microdeletions, particularly the signs and symptoms of Gorlin syndrome, result from a loss of the *PTCH1* gene. Other signs and symptoms related to 9q22.3 microdeletions probably result from the loss of additional genes in the q22.3 region. Researchers are working to determine which missing genes contribute to the other features associated with the deletion.

bladder cancer

Deletions of part or all of chromosome 9 are commonly found in bladder cancers. These chromosomal changes are seen only in cancer cells and typically occur early in tumor formation. Researchers believe that several genes that play a role in bladder cancer may be located on chromosome 9. They suspect that these genes may be tumor suppressors, which means they normally help prevent cells from growing and dividing in an uncontrolled way. Researchers are working to determine which genes, when altered or missing, are involved in the development and progression of bladder tumors.

chronic myeloid leukemia

A rearrangement (translocation) of genetic material between chromosomes 9 and 22 causes a type of cancer of blood-forming cells called chronic myeloid leukemia. This slow-growing cancer leads to an overproduction of abnormal white blood cells. Common features of the condition include excessive tiredness (fatigue), fever, weight loss, and an enlarged spleen.

The translocation involved in this condition, written as t(9;22), fuses part of the *ABL1* gene from chromosome 9 with part of the *BCR* gene from chromosome 22, creating an abnormal fusion gene called *BCR-ABL1*. The abnormal chromosome 22, containing a piece of chromosome 9 and the fusion gene, is commonly called the Philadelphia chromosome. The translocation is acquired during a person's lifetime and is present only in the abnormal blood cells. This type of genetic change, called a somatic mutation, is not inherited.

The protein produced from *BCR-ABL1* gene signals cells to continue dividing abnormally and prevents them from self-destructing, which leads to overproduction of the abnormal cells.

The Philadelphia chromosome also has been found in some cases of rapidly progressing blood cancers known as acute leukemias. It is likely that the form of blood cancer that develops is influenced by the type of blood cell that acquires the mutation and other genetic changes that occur. The presence of the Philadelphia chromosome provides a target for molecular therapies.

Kleefstra syndrome

Most people with Kleefstra syndrome, a disorder with signs and symptoms involving many parts of the body, are missing a sequence of about 1 million DNA building blocks (base pairs) on one copy of chromosome 9 in each cell. The deletion occurs near the end of the long (q) arm of the chromosome at a location designated q34.3, a region containing a gene called *EHMT1*. Some affected individuals have shorter or longer deletions in the same region.

The loss of the *EHMT1* gene from one copy of chromosome 9 in each cell is believed to be responsible for the characteristic features of Kleefstra syndrome in people with the 9q34.3 deletion. However, the loss of other genes in the same region may lead to additional health problems in some affected individuals.

The *EHMT1* gene provides instructions for making an enzyme called euchromatic histone methyltransferase 1. Histone methyltransferases are enzymes that modify proteins called histones. Histones are structural proteins that attach (bind) to DNA and give chromosomes their shape. By adding a molecule called a methyl group to histones, histone methyltransferases can turn off (suppress) the activity of certain genes, which is essential for normal development and function. A lack of euchromatic histone methyltransferase 1 enzyme impairs proper control of the activity of certain genes in many of the body's organs and tissues, resulting in the abnormalities of development and function characteristic of Kleefstra syndrome.

other cancers


Changes in the structure of chromosome 9 have been found in many types of cancer. These changes, which occur only in cells that give rise to cancer, usually involve a loss of part of the chromosome or a rearrangement of chromosomal material. For example, a loss of part of the long (q) arm of chromosome 9 has been identified in some types of brain tumor. In addition, chromosomal rearrangements that fuse the *ABL1* gene with genes other than *BCR* have been found in a small number of acute leukemias. The exact mechanisms by which these genetic changes lead to cancer are not completely understood, although it is likely that the proteins produced from them promote uncontrolled growth of cells.

other chromosomal conditions

Other changes in the structure or number of copies of chromosome 9 can have a variety of effects. Intellectual disability, delayed development, distinctive facial features, and an unusual head shape are common features. Changes to chromosome 9 include an extra piece of the chromosome in each cell (partial trisomy), a missing segment of the chromosome in each cell (partial monosomy), and a circular structure called a ring chromosome 9. A ring chromosome occurs when both ends of a broken chromosome are reunited. Rearrangements (translocations) of genetic material between chromosome 9 and other chromosomes can also lead to extra or missing chromosome segments.

Chromosome Diagram

Geneticists use diagrams called idiograms as a standard representation for chromosomes. Idiograms show a chromosome's relative size and its banding pattern, which is the characteristic pattern of dark and light bands that appears when a chromosome is stained with a chemical solution and then viewed under a microscope. These bands are used to describe the location of genes on each chromosome.



Credit: Genome Decoration Page/NCBI

Additional Information & Resources

MedlinePlus

- Encyclopedia: Chromosome
<https://medlineplus.gov/ency/article/002327.htm>

Additional NIH Resources

- National Human Genome Research Institute: Chromosome Abnormalities
<https://www.genome.gov/11508982/>

Educational Resources

- Genome News Network: Human Chromosomes 9 and 10 Are Complete (May 26, 2004)
<http://www.genomenewsnetwork.org/articles/2004/05/26/chromosomes.php>

GeneReviews

- 9q22.3 Microdeletion
<https://www.ncbi.nlm.nih.gov/books/NBK61984>
- Kleeftstra Syndrome
<https://www.ncbi.nlm.nih.gov/books/NBK47079>

Scientific Articles on PubMed

- PubMed
<https://www.ncbi.nlm.nih.gov/pubmed?term=%28Chromosomes,+Human,+Pair+9%5BMAJR%5D%29+AND+%28Chromosome+9%5BTI%5D%29+AND+english%5BIa%5D+AND+human%5Bmh%5D+AND+%22last+3600+days%22%5Bdp%5D>

Sources for This Summary

- Chereda B, Melo JV. Natural course and biology of CML. *Ann Hematol.* 2015 Apr;94 Suppl 2: S107-21. doi: 10.1007/s00277-015-2325-z. Epub 2015 Mar 27. Review.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/25814077>
- Coco S, Valdora F, Bonassi S, Scaruffi P, Stigliani S, Oberthuer A, Berthold F, Andolfo I, Servidei T, Riccardi R, Basso E, Iolascon A, Tonini GP. Chromosome 9q and 16q loss identified by genome-wide pooled-analysis are associated with tumor aggressiveness in patients with classic medulloblastoma. *OMICS.* 2011 May;15(5):273-80. doi: 10.1089/omi.2010.0103. Epub 2011 Feb 24.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/21348762>
- De Braekeleer E, Douet-Guilbert N, Rowe D, Bown N, Morel F, Berthou C, Férec C, De Braekeleer M. ABL1 fusion genes in hematological malignancies: a review. *Eur J Haematol.* 2011 May;86(5): 361-71. doi: 10.1111/j.1600-0609.2011.01586.x. Epub 2011 Mar 23. Review.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/21435002>
- Ensembl Human Map View
http://www.ensembl.org/Homo_sapiens/Location/Chromosome?chr=9;r=9:1-138394717
- Gilbert F, Kauff N. Disease genes and chromosomes: disease maps of the human genome. Chromosome 9. *Genet Test.* 2001 Summer;5(2):157-74.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/11551106>
- Hoischen A, Ehrler M, Fassunke J, Simon M, Baudis M, Landwehr C, Radlwimmer B, Lichter P, Schramm J, Becker AJ, Weber RG. Comprehensive characterization of genomic aberrations in gangliogliomas by CGH, array-based CGH and interphase FISH. *Brain Pathol.* 2008 Jul;18(3): 326-37. doi: 10.1111/j.1750-3639.2008.00122.x. Epub 2008 Mar 26.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/18371186>
- Humphray SJ, Oliver K, Hunt AR, Plumb RW, Loveland JE, Howe KL, Andrews TD, Searle S, Hunt SE, Scott CE, Jones MC, Ainscough R, Almeida JP, Ambrose KD, Ashwell RI, Babbage AK, Babbage S, Bagguley CL, Bailey J, Banerjee R, Barker DJ, Barlow KF, Bates K, Beasley H, Beasley O, Bird CP, Bray-Allen S, Brown AJ, Brown JY, Burford D, Burrill W, Burton J, Carder C, Carter NP, Chapman JC, Chen Y, Clarke G, Clark SY, Clee CM, Clegg S, Collier RE, Corby N, Crosier M, Cummings AT, Davies J, Dhami P, Dunn M, Dutta I, Dyer LW, Earthrowl ME, Faulkner L, Fleming CJ, Frankish A, Frankland JA, French L, Fricker DG, Garner P, Garnett J, Ghori J, Gilbert JG, Glison C, Grafham DV, Gribble S, Griffiths C, Griffiths-Jones S, Grocock R, Guy J, Hall RE, Hammond S, Harley JL, Harrison ES, Hart EA, Heath PD, Henderson CD, Hopkins BL, Howard PJ, Howden PJ, Huckle E, Johnson C, Johnson D, Joy AA, Kay M, Keenan S, Kershaw JK, Kimberley AM, King A, Knights A, Laird GK, Langford C, Lawlor S, Leongamornlert DA, Leversha M, Lloyd C, Lloyd DM, Lovell J, Martin S, Mashreghi-Mohammadi M, Matthews L, McLaren S, McLay KE, McMurray A, Milne S, Nickerson T, Nisbett J, Nordsiek G, Pearce AV, Peck AI, Porter KM, Pandian R, Pelan S, Phillimore B, Povey S, Ramsey Y, Rand V, Scharfe M, Sehra HK, Shownkeen R, Sims SK, Skuce CD, Smith M, Steward CA, Swarbreck D, Sycamore N, Tester J, Thorpe A, Tracey A, Tromans A, Thomas DW, Wall M, Wallis JM, West AP, Whitehead SL, Willey DL, Williams SA, Wilming L, Wray PW, Young L, Ashurst JL, Coulson A, Blöcker H, Durbin R, Sulston JE, Hubbard T, Jackson MJ, Bentley DR, Beck S, Rogers J, Dunham I. DNA sequence and analysis of human chromosome 9. *Nature.* 2004 May 27;429(6990):369-74.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/15164053>
Free article on PubMed Central: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2734081/>
- Map Viewer: Genes on Sequence
<https://www.ncbi.nlm.nih.gov/mapview/maps.cgi?ORG=human&MAPS=ideogr,ugHs,genes&CHR=9>

- Mhawech-Fauceglia P, Cheney RT, Schwaller J. Genetic alterations in urothelial bladder carcinoma: an updated review. *Cancer*. 2006 Mar 15;106(6):1205-16. Review.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/16470587>
- Muller EA, Aradhya S, Atkin JF, Carmany EP, Elliott AM, Chudley AE, Clark RD, Everman DB, Garner S, Hall BD, Herman GE, Kivuva E, Ramanathan S, Stevenson DA, Stockton DW, Hudgins L. Microdeletion 9q22.3 syndrome includes metopic craniosynostosis, hydrocephalus, macrosomia, and developmental delay. *Am J Med Genet A*. 2012 Feb;158A(2):391-9. doi: 10.1002/ajmg.a.34216. Epub 2011 Dec 21.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/22190277>
- Redon R, Baujat G, Sanlaville D, Le Merrer M, Vekemans M, Munnich A, Carter NP, Cormier-Daire V, Colleaux L. Interstitial 9q22.3 microdeletion: clinical and molecular characterisation of a newly recognised overgrowth syndrome. *Eur J Hum Genet*. 2006 Jun;14(6):759-67.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/16570072>
- Stewart DR, Kleefstra T. The chromosome 9q subtelomere deletion syndrome. *Am J Med Genet C Semin Med Genet*. 2007 Nov 15;145C(4):383-92. Review.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/17910072>
- UCSC Genome Browser: Statistics
<http://genome.cse.ucsc.edu/goldenPath/stats.html>
- Willemsen MH, Beunders G, Callaghan M, de Leeuw N, Nillesen WM, Yntema HG, van Hagen JM, Nieuwint AW, Morrison N, Keijzers-Vloet ST, Hoischen A, Brunner HG, Tolmie J, Kleefstra T. Familial Kleefstra syndrome due to maternal somatic mosaicism for interstitial 9q34.3 microdeletions. *Clin Genet*. 2011 Jul;80(1):31-8. doi: 10.1111/j.1399-0004.2010.01607.x. Epub 2011 Jan 10.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/21204793>
- Wolff EM, Liang G, Jones PA. Mechanisms of Disease: genetic and epigenetic alterations that drive bladder cancer. *Nat Clin Pract Urol*. 2005 Oct;2(10):502-10. Review.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/16474624>
- Yatsenko SA, Brundage EK, Roney EK, Cheung SW, Chinault AC, Lupski JR. Molecular mechanisms for subtelomeric rearrangements associated with the 9q34.3 microdeletion syndrome. *Hum Mol Genet*. 2009 Jun 1;18(11):1924-36. doi: 10.1093/hmg/ddp114. Epub 2009 Mar 17.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/19293338>
Free article on PubMed Central: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2678925/>

Reprinted from Genetics Home Reference:
<https://ghr.nlm.nih.gov/chromosome/9.pdf>

Reviewed: September 2016

Published: March 21, 2017

Lister Hill National Center for Biomedical Communications
U.S. National Library of Medicine
National Institutes of Health
Department of Health & Human Services